



## REGIONAL FEATURES OF THE COURSE OF SURGICAL INFECTION AROUND THE BUKHARA REGION

Nurboboyev Adhambek Uyg'unovich

Bukhara state medical institute

<https://doi.org/10.5281/zenodo.11634623>

### ARTICLE INFO

Qabul qilindi: 05-June 2024 yil

Ma'qullandi: 10- June 2024 yil

Nashr qilindi: 15- June 2024 yil

### KEYWORDS

*regional surgery, infection, sensitive to antibiotics, the resistance*

### ABSTRACT

*The Bukhara region of the republic of Uzbekistan at the who received the western area, which produces a purulent surgical diseases of soft tissues in the microflora epidemiological study was conducted, including covers. The materials obtained on the basis of the type of surgical infections resistant to antibiotics and the epidemiological characteristics of the microflora were coming, and of the bukhara region of them atnu he who received the western region of growth in the scale identified. Soft and purulent surgical diseases of pathogenic microflora of the research came from the specific characteristics of the regional differences shows..*

**The actual ministry.** Purulent-surgical treatment of diseases of the modern surgeon remains a pressing problem. According to the opinion of many authors have been at the hospital all of purulent surgical diseases of surgical pathology requiring treatment from 30% to 40% (0). This indicator, our ma'lumotlarimizga according outpatient surgical diseases in the analysis of purulent grow more.

Infection surgery, including postoperative purulent infections of the problem of the cart in June 2015 and September 2016 was considered within the framework of the un general assembly in this issue 71-considered in the session, was attended by experts from all countries of the world. Postoperative wound infections are a common complication following surgery, characterized by complex and multifactorial pathophysiology. Clinicians are crucial in identifying and managing modifiable risk factors for postoperative wound infections during the perioperative phase. Comprehensive preoperative assessment and management are essential, necessitating collaboration among nursing, anesthesia, and surgical teams to identify and manage risk factors, thereby ensuring effective patient counseling. During the intraoperative phase, maintaining sterility and cleanliness in the operating room environment is crucial, as it directly influences patient recovery and infection rates intraoperatively and postoperatively.

Incisional infections are usually noticeable, and any systemic symptoms following surgery should raise concerns about postoperative complications. However, similar symptoms may also stem from unrelated causes, such as cellulitis, allergic reactions, or urinary tract infections and pneumonia. Various factors contribute to susceptibility to infection, and

diagnosis primarily relies on clinical evaluation, although wound cultures and imaging may be necessary in some instances. This activity provides detailed information on postoperative wound infections, including their etiology, epidemiology, pathophysiology, common presentations, and evaluation and treatment options. This activity also highlights the crucial role of the interprofessional healthcare team and the collaborative approach between surgical and non-surgical clinicians to ensure the best possible patient outcomes.

Purulent diseases of infection associated with surgery have a tendency to grow around the world and the world of medicine's most pressing problem was noted that the fact that the wound was also developed strategies to fight against purulent infections and to prevent them. As noted, this microflora of taking the specific features of each region for the study of the sensitivity of the microflora play an important role in. In particular, we paid attention to increase the sensitivity of the microflora and the way to avoid them.

**Research purpose:** region Bukhara purulent surgical infection study the characteristics of regional the night.

**Research materials and methods.**

Region bukhara atnu he who received the western territory, Karakul district of 2023 in the year of purulent surgery, including infection treatment covers the analysis of the results was. We are based outpatient treatment for purulent surgical diseases of soft tissues 529 records we have analyzed outpatient units. Quantitative and qualitative bacteriological studies and the responsiveness of microflora to antibiotics in all patients studied.

Microbiological analysis of wound infection, as well as the qualitative and quantitative assessment of sensitivity to various antibiotics was carried out through implementation sekresiya collection from the wound.

**The method for determining resistance of microflora.**

The drive resistance of microflora balances diffusion method (DDhe) to identify its essence is as follows:

Method of application: to determine the sensitivity of ddhe 's dissolve into the environment from paper disks impregnated with abp feed them if fertilization on the growth of microorganisms on the surfacet to prevent is based on its ability to.

**Get the results into account.**

After completion of incubation, the occurrence of the dark mat of the vessel is put in reverse, then to them the light of 45 ° at an angle falls (light is reflected). The delay is measured with an accuracy of 1 mm zone diameter of growth, it is preferable to use or kaliper kaliperdan.


Growth to prevent visible growth of the zone to the zone of complete suppression of the measurement need to pay attention to. Growth to prevent light conditions detected in very small colonies in the zone to zoom in or to only on special zone and you should not give attention to at the edge of the covering almost not felt. The results of the sensitivity of detection of staphylococcus oksatsilinga, taking into account the zone of growth of small colonies identified in most of the exact inhibisyon when you need to take is no exception. microflora populations of microorganisms, or the presence of a strange geterorezistentligi this colony should determine the sensitivity of these strains to identify microorganisms and repeat the executive. Sar meat proteus strainsto determine the sensitivity of dd ng my growth to prevent the thin film can be covered with a veil similar to that of the zone, border

zone and not to interfere with the results recorded in this character does not take into account.

Sulfanilamidlar are sensitive to inhibit the growth zone and their interaction in combination with trimetoprim to determine the limit of 80% of growth to prevent the level should be taken into account. The reason for this is that these microorganisms under the influence of drugs against a full understanding of the reproduction of before the end of the growth cycle until 1-2 can be.


**Results and their discussion.**

That covers the analysis of the results of the study, including analysis, Karakul district 's total population number 169 669 students. Purulent diseases of soft tissues in the year of 2023 to universal central polyclinic surgical 529 people (the population of the region of 0,2 percent) patient contact, that 472 students of the patient is based outpatient treatment. Of them, 348 dollar (73,7%) purulent wounds of soft tissues, 20 dollars, (4,2%) or gavin foot-whose hands flegmonasi, 4 of them in (0,8%) furunkul, 97 in them (20,5%) and purulent wounds of the fingers, and 3 of them in (0,6%) abses with.



**Rice. The patient distribution by type 1 of the disease.**

Identify the following features of the results of bacteriological research: st. aureus 360 (61%), st.Epidermidis 60 (10,1%), Streptococcus 120 (20,3%), ps.aerogenaza 50 (8,6%) strains detected in 2-picture.



2-picture. Distribution of patients according to the type of microflora.

As noted above, the microflora of purulent see sensitive to antibiotics was also studied. Identify, **least tree-cutting Aureus** was sensitive to the following antibiotics: sefozalin, kanamitsin 21,6%, gentamits may be 47,5%, ceftriaxone, amoksiklav 22,7%, levfloksatsin 48,3%, eritromitwith 22,2% and 1 table. **Streptococcal** microflora at theoksiklav 46,6%, sefapirozin 54,1%, gentamicin, linkomitsin 49,1%, kanamitsin 45,8% s like antibioticsdetected infringement ezuvchi 2-table. **st.Epidermidis** s ateftriakson 88,3% and klovanir 85% sensitivity detected 3-table. **Ps.aerogenaza** the following antibiotics to determine susceptibilty levoroks 100%, sefadoksim 100% 4-table.

Each of the type of microflora to antibiotics sensitive are listed in the following table:

1 Picture.

Sensitivity the least tree-cutting Aureusdan antibiotics (n=360)

No	Antibiotic is the name of the	Sensitivity of the number of	Percentage (%)
1	Kanamisin	78	21,6%
2	Tsefozolin	78	21,6%
3	Gentamicin	171	47,5%
4	Tseftriakson	82	22,7%
5	perks of amoiklav	82	22,7%
6	Levofloksasin	174	48,3%
7	Eritromisin	80	22,2%

2 Picture.

Streptococcus antibiotics sensitivity (n=120)

No	Antibiotic is the name of the	Sensitivity of the number of	Percentage (%)
----	-------------------------------	------------------------------	----------------

1	Amoksiklav	56	46,6%
2	Tsefapirozin	65	54,1%
3	Gentamicin	59	49,1%
4	Linkomisin	59	49,1%
5	Kanamisin	55	45,8%

**3 Picture.**

**st.Epidermidis 's antibiotics sensitivity (n=60)**

No	Antibiotic is the name of the	Sensitivity of the number of	Percentage (%)
1	Tseftriakson	53	88,3%
2	Klovanir	51	85%

**4 Picture.**

**Ps.aerogenaza 's antibiotiklarga sensitivity (n=50)**

No	Antibiotic is the name of the	Sensitivity of the number of	Percentage (%)
1	Levoroks	50	100%
2	Tsefodoksim	50	100%

Thus, our research to identify the following interesting thoughts - learning we bukhara province of the south western territory, the frequency of purulent surgical diseases, pathogenic microflora of them has specific properties and the type of resistance to various antibiotics. purulent diseases of soft tissues and outpatient diagnosis and treatment of conditions that should be taken into account.

**Summary:**

1. Pathogenic microflora of purulent surgical diseases of soft tissues in each region come from specific features.
2. Each is a region of microflora resistance properties separately study should.
3. Every a microflora of different antibiotic sensitivity own specific characteristics has.

**References:**

1. Khalmatova, B. T., Tashmatova, G. A., & Mirsalikhova, N. K. (2021). Modern methods for diagnosing the function of external respiration in children with bronchial asthma. *ACADEMICIA: An International Multidisciplinary Research Journal*, 11(4), 844-847.
2. Abdullayeva, M. (2019). Clinical efficacy of montelukast (l-montus kid®) in the control of Mild persistent bronchial asthma in children. *Journal of Critical Reviews*, 7(5), 2020.
3. Юлдашева, Ф. У., & Тошматова, Г. А. (2023). ОКАЗАНИЕ МЕДИКО-СОЦИАЛЬНОЙ ПОМОЩИ НАСЕЛЕНИЮ.

4. Khalmatova, B. T., & Tashmatova, G. A. (2021). Influence Of Ecological Factors On The Development And Progress Of Bronchial Asthma In Children. *European Journal of Molecular & Clinical Medicine*, 7(8), 4374-4377.
5. Mirrahimova, M. H., Khalmatova, B. T., & Tashmatova, G. A. (2019). Bronchial asthma in children: a modern view of the problem. *Toshkent tibbiyot akademiyasi axborotnomasi*, (1), 31-34.
6. Миррахимова, М. Х., Халматова, Б. Т., & Тошматова, Г. А. (2019). Бронхиальная астма у детей: современный взгляд на проблему.
7. Шамсиева, Э. Р., & Ташматова, Г. А. (2022). Деструктивно-воспалительная патология суставов у детей до 16 лет (Doctoral dissertation, England).
8. Tashmatova, G. A., & Khalmatova, B. T. (2021). Bronchial asthma in children during the covid-19 pandemic: a feature of the course.
9. Mirrahimova, M. H., & Khalmatova, B. T. (2019). Prevalence and risk factors of bronchial asthma in children living in the industrial zones of Tashkent region. *Central Asian Journal of Medicine*, 2018(4), 67-76.
10. Maktuba, T., Dildoraxon, N., Elnora, D., Dilnoza, S., & Nurbek, A. (2020). 14-3-3 (eta) protein as a new biomarker in rheumatoid arthritis. *International Journal of Advanced Science and Technology*, 29(5), 1871-1877.
11. Джураева, Э. Р., & Ганиева, Н. А. (2016). Современные подходы к лечению рецидивирующего полихондрита. *Вопросы организации и информатизации здравоохранения*, (S), 116-117.
12. Ganiyeva, N. A., Djurayeva, E. R., Abduazizova, N. X., Aripova, N. A., Kasimova, M. B., Axmedova, N. A., & Bekenova, G. T. (2023). ASSESSMENT OF CLINICAL AND DIAGNOSTIC ASPECTS OF ATHEROSCLEROSIS IN PATIENTS WITH SYSTEMIC SCLERODERMA. *湖南大学学报 (自然科学版)*, 50(07).
13. Abduazizova, N. X., Nabieva, D. A., & Sultanov, J. A. (2023). Lipid profile in patients with rheumatoid arthritis on the background of basic treatment. *Asian journal of pharmaceutical and biological research*, 12(2).
14. Djurayeva, E. R., & Normuradov, A. D. (2023). Importance of coronavirus infection (covid-19) in the development of systemic vasculitis. *Advanced Performance Materials*, 2(1).
15. Pulatova, S., Nabiyeva, D., Abduazizova, N., Mukhammadiyeva, S., & Agzamova, G. (2022). Clinical and pathogenetic values of disorders of mineral metabolism in ankylosing spondylitis.
16. Ganiyeva, N. A., Rizamukhamedova, M. Z., Nabiyeva, D. A., & Aripova, N. A. (2021). Clinic-Diagnostic Aspects of Modern Biomarkers of Early Atherosclerosis and Fibrotic activity of Systemic Scleroderma. *Asian Journal of Medical Principles and Clinical Practice*, 4(3), 1-13.
17. Aripov, A. N., Aripov, O. A., Akhundjanova, L. L., Nabiev, A. U., Nabieva, D. A., & Khamroev, T. T. (2022). Study the antifibrous efficacy of plant proanthocyanidin in rats with chronic heliotrine liver damage.
18. Tashpulatova, M. M., Nabiyeva, D. A., Djurayeva, E. R., & Akhmedova, N. A. (2021). Diagnostic Significance of 14-3-3  $\eta$  (Eta) Protein and MRI of Joints in Early Stage of Rheumatoid Arthritis.

19. Nabieva, D. A. (2017). AN Aripov The detection of proteomic markers and immunologic profile and their relationship with metabolic parameters in patients with gout. *Klinicheskaia Laboratornaia Diagnostika*, (62), 8.
20. Джураева, Э. Р., & Ганиева, Н. А. (2016). Современные подходы к лечению рецидивирующего полихондрита. *Вопросы организации и информатизации здравоохранения*, (S), 116-117.
21. Ganiyeva, N. A., Djurayeva, E. R., Abduazizova, N. X., Aripova, N. A., Kasimova, M. B., Axmedova, N. A., & Bekenova, G. T. (2023). ASSESSMENT OF CLINICAL AND DIAGNOSTIC ASPECTS OF ATHEROSCLEROSIS IN PATIENTS WITH SYSTEMIC SCLERODERMA. *湖南大学学报 (自然科学版)*, 50(07).
22. Aripova, N. A., Djurayeva, E. R., Abduazizova, N. X., Berdiyeva, D. U., Ganiyeva, N. A., & Ziyayeva, F. K. (2023). The role of pro-inflammatory cytokines in various types of systemic scleroderma. *Seybold Rep. J*, 18(06), 1469-1476.
23. Ганиева, Н. А., Джураева, Э. Р., & Арипова, Н. А. (2022). Комбинированная терапия синдрома Рейно при системной склеродермии (Doctoral dissertation).



INNOVATIVE  
ACADEMY