



## SELECTION OF EARLY MATURING AND HIGH YIELDING LINES OF DURUM WHEAT FOR IRRIGATED AREAS

Dilmurodov Sherzod Dilmurodovich <sup>1</sup>  
Abdimajidov Jaloliddin Raxmatullaevich <sup>2</sup>  
Shodiyev Sherzod Shomiljonovich <sup>3</sup>  
Boysunov Nurzod Bekmurodovich <sup>4</sup>  
Kayumov Norboy Shakirjanovich <sup>5</sup>

<sup>1</sup> Kashkadarya branch of the Resspikech Institute for Grain and Leguminous crops  
Doctor of Philosophy (PhD) in Agricultural Science, Senior scientific researcher  
<sup>2,3,4,5</sup> PhD students. Junior scientific researchers

<https://doi.org/10.5281/zenodo.4733079>

### ARTICLE INFO

Received: 25<sup>th</sup> April 2021  
Accepted: 27<sup>th</sup> April 2021  
Online: 29<sup>th</sup> April 2021

### KEY WORDS

*durum wheat, variety, lines, early maturity, grain yield.*

### ABSTRACT

*Currently, the import of large quantities of pasta and confectionery products from foreign countries, which are necessary for the needs of the population, affects the state budget. The quality of grain grown often does not fully meet the requirements of the bakery, pasta and confectionery industries. One of the most pressing issues in agriculture is the creation of new high-yielding, fast-maturing, dormant and disease-resistant varieties in the cultivation of durum wheat. The article is based on the study of new forms of hybrid lines to create early-maturing and yielding varieties of durum wheat.*

The existing wheat varieties created by breeders are required to maintain their high yield, disease and pest resistance in different climatic conditions of the year, as well as high quality indicators in the competitive varietal testing nursery of durum wheat [2, 10, 21, 27, 34].

The high protein content and high gluten content of durum wheat ensure that pasta made from it is yellow-gold or lemon-colored and does not crumble during cooking [6, 9, 16, 25, 35].

Durum wheat is the only and irreplaceable raw material for the production of pasta and wheat bran. It has original properties compared to other wheat products: high transportability, taste and nutritional quality even when stored in a dry place for a long time, ease of cooking in a minimum time, relatively low cost, versatility, dietary product for patients [7, 12, 20, 22, 30].

Depending on the natural-climatic conditions of each region, it is necessary to create new varieties with a new genotype, high yields, alternate yields and good quality [1, 13, 17, 26, 33].

Development and introduction of new varieties of durum wheat requires high heat, drought tolerance and high yield elements [4, 11, 15, 23, 31].

Increasing wheat production to meet protein demand, improving the structure of arable land, efficient use of mineral and organic fertilizers, the use of high agronomic techniques, can be achieved with varieties with high protein content [3, 8, 19, 24, 29].

Durum wheat varieties are less resistant to drought, high temperatures, yellow and brown rust and black spot disease than soft wheat varieties, and are



distinguished by the fact that the grains in the ears do not spill when ripe [5, 14, 18, 28, 32].

20 durum wheat varieties and lines were selected as the object of study. Krupinka variety, which is planted for irrigated areas, was taken as a standard variety. In Experiment 3, the crop area was planted at 10 m<sup>2</sup>.

Field experiment was conducted in the Karshi experimental field of the Kashkadarya branch of the Scientific Research Institute of Cereals and Legumes. Experimental placement and experimentation were carried out according to the method of phenological observation, calculation and analysis (All-Union Institute of Botany VIR, 1984) and biometric analysis according to the method of the State Variety Testing Commission of Agricultural Crops (1985, 1989). Statistical analyzes were performed based on the method of B.A. Dospikhov (1985). The coefficient of interdependence of characters is calculated on the basis of the method of P.V. Terentev (1959). The field experimental scheme in the study was based on the Complete block design and Alpha lattice design of the Genstat 3 program.

The experiment was sown on October 22 at the expense of 5 million germinated seeds. Irrigation with seed water was carried out to fully recover the planted varieties. According to the results of phenological

observations, germination of varieties and lines occurred from 1 to 2 November. The accumulation phase was observed on November 30 in 3 lines, and on December 1-3 in the remaining 17 cultivars and lines. The piping phase lasted from February 27 to March 6.

The maturing period of the variety and lines is the most favorable stage for the separation of early maturing lines. In the experiment, it was noted that the grazing phase of the varietal lines began a little earlier, depending on the weather, and lasted from 4 to 16 April. While the standard Krupinka variety entered the germination phase on April 9, 9 lines entered the early stage, 8 lines entered the late stage, and 2 lines entered the germination phase at the same time as the standard variety.

The maturing period was from May 25 to June 4, and the growth period was found to be 204 to 215 days. While the standard Krupinka variety entered the full maturing phase on May 29, 16 lines were found to ripen earlier than the standard variety. KR17\_DurumYT\_Entry-18, KR17\_DurumAgro\_Entry-3, KR17\_DurumYT\_Entry-15, KR17\_DurumYT\_Entry-16 lines were assessed as early maturing and were found to have entered the full maturing phase on 25 May.

Table 1

### Results of phenological observation and biometric measurements of varieties and lines (Karshi, 2018).

№	Name	Heading date	Days to heading date	Maturity date	Days to maturity date	Plant height, cm	Peduncle length, cm	Spike length, cm	Number of spikelets of per spike	Grain yield, c/ha
1	KR17_DurumAgro_Entry-3	8 апр	158	25 май	205	86,0	39,3	6,3	17	75,4
2	KR17_DurumAgro_Entry-7	9 апр	159	28 май	207	91,3	38,7	5,3	17	85,8
3	KR17_DurumAgro_Entry-16	8 апр	157	26 май	205	89,7	43,0	6,0	17	83,2
4	KR17_DurumAgro_Entry-18	13 апр	163	28 май	209	96,0	39,3	6,0	17	75,0



5	KR17_DurumAgro_Entry-19	10 апр	160	31 май	211	90,7	36,3	5,7	16	82,0
6	KR17_DurumAgro_Entry-20	9 апр	159	27 май	207	92,3	36,3	6,7	17	77,2
7	KR17_DurumYT_Entry-2	7 апр	157	26 май	206	89,0	35,0	6,0	18	87,3
8	KR17_DurumYT_Entry-3	5 апр	155	26 май	206	81,7	35,7	5,3	15	85,1
9	KR17_DurumYT_Entry-7	5 апр	155	26 май	206	94,7	36,7	7,0	18	73,9
10	KR17_DurumYT_Entry-14	10 апр	160	27 май	206	102,7	43,3	8,3	19	82,7
11	KR17_DurumYT_Entry-15	4 апр	153	25 май	204	88,0	44,3	6,3	17	62,3
12	KR17_DurumYT_Entry-16	5 апр	155	25 май	205	76,0	37,0	5,7	15	62,5
13	KR17_DurumYT_Entry-18	4 апр	154	25 май	204	95,3	41,0	6,7	17	74,8
14	KR17_DurumYT_Entry-19	7 апр	156	26 май	205	79,7	38,0	5,3	15	81,8
15	KRDW17-1	13 апр	162	4 июн	215	96,0	34,3	7,3	21	81,5
16	KRDW17-2	16 апр	165	1 июн	212	90,0	32,7	6,7	17	71,7
17	KRDW17-3	11 апр	162	26 май	206	86,7	36,7	5,7	16	69,0
18	KRDW17-4	10 апр	160	28 май	208	82,3	37,3	5,3	17	74,8
19	KRDW17-5	10 апр	160	28 май	208	91,7	40,0	7,7	19	78,0
20	Krupinka (андоза)	9 апр	159	29 май	208	94,0	32,3	7,3	20	73,4
	<b>Mean</b>	<b>8 апр</b>	<b>158</b>	<b>27 май</b>	<b>207</b>	<b>89,7</b>	<b>37,9</b>	<b>6,3</b>	<b>17,1</b>	<b>76,9</b>
	<b>Maximum</b>	<b>16 апр</b>	<b>165</b>	<b>4 июн</b>	<b>215</b>	<b>102,7</b>	<b>44,3</b>	<b>8,3</b>	<b>20,7</b>	<b>87,3</b>
	<b>Minimum</b>	<b>4 апр</b>	<b>153</b>	<b>25 май</b>	<b>204</b>	<b>76,0</b>	<b>32,3</b>	<b>5,3</b>	<b>14,7</b>	<b>62,3</b>
	<b>LSD<sub>0,5</sub></b>		<b>3,49</b>		<b>2,01</b>	<b>3,38</b>	<b>2,83</b>			<b>1,02</b>
	<b>LSD<sub>0,5</sub> %</b>		<b>2,2</b>		<b>1,0</b>	<b>3,8</b>	<b>7,5</b>			<b>1,3</b>
	<b>CV%</b>		<b>1,3</b>		<b>0,6</b>	<b>2,2</b>	<b>4,4</b>			<b>0,8</b>

The height of the varieties studied in the study has a significant impact on high yields. Stunted varieties have a sharp decline in yield and grain quality in drought conditions. Therefore, the choice of varieties with long plant height in irrigated areas is one of the main features.

According to the results of the study, the height of the varieties ranged from 76.0 to 102.7 cm. It was found that the number of short-stemmed lines with a plant height of less than 80 cm was 2. It was noted that the plant height of the remaining cultivars and lines ranged from 90 to 102.7 cm.

When the relationship between plant height and yield was studied, it was found that there was a positive correlation between  $r = 0.25$ . From the above, it can be seen that the height of the plant had a great influence on the high yield of the elements.

In order to determine the yield of varieties, harvesting was carried out on a

special selection combine. The average yield of returns was calculated and the yield of the cultivars and lines studied ranged from 62.3 to 87.3 ts / ha. In the experiment, it was found that the yield of Krupinka variety, taken as a standard variety, was 73.4 ts / ha, and the yield of 15 lines was higher than the standard variety. According to the results, the KR17\_DurumYT\_Entry-2 ridge showed a high yield of 87.3 ts / ha, the KR17\_DurumAgro\_Entry-7 ridge 85.8 ts / ha, the KR17\_DurumYT\_Entry-3 ridge 85.1 ts / ha, the KRDW17-1 ridge 81.5 ts / ha. .

In conclusion, it should be noted that from the studied durum wheat lines (KR17\_DurumYT\_Entry-18, KR17\_DurumYT\_Entry-3, KR17\_DurumYT\_Entry-19, KR17\_DurumYT\_Entry-2, KR17\_DurumAchro\_Entry) were transferred to Agroecological yield trial.



## REFERENCES:

1. Juraev D. T. et al. To study the heat resistance features of bread wheat varieties and species for the southern regions of the republic of Uzbekistan //European Journal of Molecular & Clinical Medicine. – 2020. – Т. 7. – №. 2. – С. 2254-2270.
2. Дилмуродов Ш. Д. Подбор исходного материала для селекции пшеницы озимой мягкой для условий Узбекистана на основе изучения хозяйственно ценных характеристик //Аграрная наука. – 2018. – №. 2. – С. 58-61.
3. Мейлиев Т. Х., Дилмуродов Ш. Д. Рост и развитие, урожайность и устойчивость к желтой ржавчине сортов в питомнике отбора продуктивных сортов //Приоритетные направления развития науки и образования. – 2019. – С. 130-133.
4. Juraev D. T. et al. Influence of hot dry winds on productivity elements of wheat crop observed in southern regions of the republic of uzbekistan //International jurnal of applied and pure science and agriculture. ISSN. – 2017. – С. 2394-5532.
5. Sh K. N. et al. Selection of spikely bread wheat lines based on studying the time of development //INTERNATIONAL SCIENTIFIC AND TECHNICAL JOURNAL “INNOVATION TECHNICAL AND TECHNOLOGY”. – 2020. – Т. 1. – №. 2. – С. 69-71.
6. Дилмуродов Ш. Д., Бойсунов Н. Б. Юмшоқ буғдой дурагай тизмаларида хосилдорликнинг биометрик кўрсаткичларга боғлиқлиги //Инновацион технологиялар. – 2020. – №. 2 (38).
7. Fayzullayev A. Z. et al. SELECTION OF HIGH-YIELDING AND HIGH-QUALITY LINES OF BREAD WHEAT //INTERNATIONAL SCIENTIFIC AND TECHNICAL JOURNAL “INNOVATION TECHNICAL AND TECHNOLOGY”. – 2020. – Т. 1. – №. 3. – С. 10-14.
8. Odirovich J. F., Anvarovich A. O., Dilmurodovich D. S. VALUABLE PROPERTIES AFFECTING THE HIGH-YIELD ELEMENTS OF DURUM WHEAT //INTERNATIONAL JOURNAL OF DISCOURSE ON INNOVATION, INTEGRATION AND EDUCATION. – 2020. – Т. 1. – №. 2. – С. 37-41.
9. Дилмуродов Ш. Д. Юмшоқ буғдойнинг маҳаллий маҳсулдор тизмалари селекцияси //Life Sciences and Agriculture. – 2020. – №. 1.
10. Juraev D. T. et al. Heritability of Valuable Economic Traits in the Hybrid Generations of Bread Wheat //Annals of the Romanian Society for Cell Biology. – 2021. – С. 2008-2019.
11. Дилмуродов Ш. Д., Жабаров Ф. О. Селекция высокоурожайных линий озимой твёрдой пшеницы с высоким качеством зерна //Молодой ученый. – 2019. – №. 31. – С. 34-38.
12. Дилмуродов Ш. Д., Бойсунов Н. Б. Отбор продуктивных линий мягких пшениц из гибридного питомника в условиях южного региона Республики Узбекистан //World Science: Problems and Innovations. – 2018. – С. 58-60.
13. Khushvaktovich M. A., Dilmurodovich D. S. THE CHOICE OF SPIKELY MATURING LINES OF SPRING BREAD WHEAT FOR IRRIGATED AREAS //НАУКА, ОБРАЗОВАНИЕ, ОБЩЕСТВО: АКТУАЛЬНЫЕ ВОПРОСЫ. – 2021. – С. 30.
14. Дилмуродов Ш. Д., Бойсунов Н. Б. Селекция местных гибридных линий мягкой пшеницы на юге Республики Узбекистан //Сборник материалов. – 2018. – С. 113-119.
15. Dilmurodov S. Some valuable properties in evaluating the productivity of bread wheat lines //INTERNATIONAL SCIENTIFIC AND TECHNICAL JOURNAL “INNOVATION TECHNICAL AND TECHNOLOGY”. – 2020. – Т. 1. – №. 1. – С. 60-62.
16. Дилмуродов Ш. Д., Зиядуллаев З. Ф. Юмшоқ буғдойда ўтказилган оддий ва мураккаб дурагайлаш ишлари натижалари //Life Sciences and Agriculture. – 2020. – №. 2.
17. Дилмуродов Ш. Д., Бойсунов Н. Б. Рақобатли нав синаш кўчатзориди юмшоқ буғдойнинг биометрик кўрсаткичларини ўрганиш //Life Sciences and Agriculture. – 2020. – №. 1.
18. Дилмуродов Ш. Д. и др. Гибридизация в различном направлении и создание гибридного поколения мягкой пшеницы //Инновационное развитие науки и образования. – 2018. – С. 74-77.



19. Дилмуродов Ш. Д., Зиядуллаев З. Ф. Selection of spikely and productive lines in preliminary yield trial of bread wheat //INTERNATIONAL SCIENTIFIC AND TECHNICAL JOURNAL "INNOVATION TECHNICAL AND TECHNOLOGY". – 2020. – Т. 1. – №. 1. – С. 55-59.
20. Дилмуродов Ш. Д. ЦЕННЫЕ СВОЙСТВА, ВЛИЯЮЩИЕ НА ВЫСОКОУРОЖАЙНЫЕ ЭЛЕМЕНТЫ МЯГКОЙ ПШЕНИЦЫ //ББК. – 2020. – Т. 60. – С. 38.
21. Dilmurodovich D. S., Bekmurodovich B. N., Shakirjonovich K. N. WINTER BREAD WHEAT GRAIN QUALITY DEPENDS ON DIFFERENT SOIL-CLIMATE CONDITIONS //INTERNATIONAL JOURNAL OF DISCOURSE ON INNOVATION, INTEGRATION AND EDUCATION. – 2020. – Т. 1. – №. 5. – С. 377-380.
22. Жураев Д. Т., Дилмуродов Ш. Д. Юмшок бутдойнинг бошоклаш-пишиш даврида иссиқликнинг таъсири //Life Sciences and Agriculture. – 2020. – №. 2-2.
23. Жураев Д. Т. и др. Влияние суховеев, наблюдаемых в южных регионах республики узбекистан, на продуктивные элементы мягкой пшеницы //Путь науки. – 2017. – №. 2. – С. 84-92.
24. Dilmurodovich D. S., Shakirjanovich K. N. ANALYSIS OF YIELD AND GRAIN QUALITY TRIATS IN THEADVANCED YIELD TRIAL OF WINTER BREAD WHEAT //Euro-Asia Conferences. – 2021. – Т. 1. – №. 1. – С. 550-555.
25. Dilmurodovich D. S. et al. Analysis of yield and yield components traits in the advanced yield trial of winter bread wheat //INTERNATIONAL JOURNAL OF DISCOURSE ON INNOVATION, INTEGRATION AND EDUCATION. – 2021. – Т. 2. – №. 1. – С. 64-68.
26. Дилмуродов Ш. Д., Каюмов Н. Ш. ОЦЕНКА ПРОДУКТИВНЫХ ПОКАЗАТЕЛЕЙ ЛИНИЙ МЯГКОЙ ПШЕНИЦЫ //Вестник науки и образования. – 2020. – №. 17-1 (95).
27. Dilmurodov S. D., Toshmetova F. N., Fayzullayeva D. SELECTION OF HIGH-QUALITY DONOR VARIETIES OF BREAD WHEAT FOR HYBRIDIZATION //МОЛОДЫЕ УЧЁНЫЕ РОССИИ. – 2020. – С. 55-58.
28. Дилмуродов Ш. Д., Каюмов Н. Ш., Бойсунов Н. Б. ЗНАЧЕНИЕ БИОМЕТРИЧЕСКИХ И ПРОДУКТИВНЫХ ПОКАЗАТЕЛЕЙ ПРИ СОЗДАНИИ ПШЕНИЦЫ С КОМПЛЕКСОМ ЦЕННЫХ СВОЙСТВ //Life Sciences and Agriculture. – 2020. – №. 2-3.
29. DILMURODOVICH D. S. et al. Productivity, quality and technological characteristics of bread wheat (*Triticum aestivum* L.) variety and lines for the southern regions of the Republic of Uzbekistan //Plant cell biotechnology and molecular biology. – 2021. – С. 63-74.
30. Dilmurodov S. D., Tukhtayeva U. A. SELECTION OF HIGH-YIELDING AND GRAIN-QUALITY DONORS OF WINTER BREAD WHEAT FOR IRRIGATED AREAS //НАУКА И ОБРАЗОВАНИЕ: СОХРАНЯЯ ПРОШЛОЕ, СОЗДАЁМ БУДУЩЕЕ. – 2020. – С. 92-95.
31. Хазраткулова Ш. У., Дилмуродов Ш. Д. Оценка жароустойчивости сортов и линий мягкой пшеницы //Наука и образование сегодня. – 2019. – №. 9 (44).
32. Dilmurodovich D. S. et al. STUDY OF MORPHO-BIOLOGICAL PROPERTIES AND RESISTANCE TO YELLOW RUST DISEASE OF NEW LINES OF WINTER BREAD WHEAT //InterConf. – 2021.
33. Shakirjanovich K. N., Dilmurodovich D. S. Analysis of yield and protein content of drought-resistant chickpea lines for rainfed areas //INTERNATIONAL JOURNAL OF DISCOURSE ON INNOVATION, INTEGRATION AND EDUCATION. – 2021. – Т. 2. – №. 1. – С. 108-111.
34. Dilmurodovich D. S., Nasirulloevna T. F. SELECTION OF SPIKELY MATURITY LINES IN AGROECOLOGICAL YIELD TRIAL OF BREAD WHEAT //НАУКА, ОБРАЗОВАНИЕ, ИННОВАЦИИ: АКТУАЛЬНЫЕ ВОПРОСЫ И. – С. 41.
35. Дилмуродов Ш. Д. и др. Сувсизликка чидамли кузги юмшок буғдой нав ва тизмаларининг баъзи қимматли хусусиятларини баҳолаш //Молодой ученый. – 2020. – №. 34. – С. 158-161.