



## PEDIATRIC HELMINTHIASIS AND THEIR INDICATIONS

**Djuraeva Muhabbat Ergashovna., Almardonova Zulxumor Jalilovna**

Assistants of Termez branch of Tashkent Medical Academy

<https://doi.org/10.5281/zenodo.7225374>

### ARTICLE INFO

Received: 03<sup>rd</sup> October 2022

Accepted: 10<sup>th</sup> October 2022

Online: 19<sup>th</sup> October 2022

### KEY WORDS

*healthy lifestyle, pathogens weaken, children's helminthosis, children's disability, acquired disability.*

### ABSTRACT

*This article presents information and conclusions about helminths in children in Surkhandarya region of Uzbekistan. According to the research results, in Surkhandarya province 44% of children are infected with enterobiasis and 23% with ascariasis, and the reasons for this are explained.*

Helminthiasis is one of the most common diseases in the world, accounting for a significant proportion of common diseases and having a negative impact on the socio-economic indicators of society. Even though 80% of cases are registered among children, it remains a constant problem among the population at this age. Pathogens weaken children's immune system and cause the development of various allergic diseases, including dermatitis, asthmatic bronchitis, rhinitis and metabolic disorders, various intestinal diseases, physical and mental weakness and many other diseases, as well as acquired disorders. It should be noted that children have high morbidity rates due to close contact with contaminated toys, household appliances and external environmental factors (soil, open water pool).

Currently, the incidence of enterobiasis (ostrich), hymenolepidosis (tapeworm), and ascariasis (ascarida) in children is 80% higher than in other types

of helminths. In children's institutions, this figure is 90%. In particular, in Surkhandarya region, in kindergartens located in Termez, Denau, Shurchi, Altynsay, Boysun, Jarkurgan and Termez districts, disinfection of children's toys is not at the necessary level, so helminthiasis is more common among children of child age. Percentage of children with helminthiasis in these districts and town of Termez is increasing day by day and number of children with disabilities is increasing.

The aim of the research. Scientific substantiation of characteristics of child helminthiasis in surkhandarya region (by the example of individual districts).

Research material and methods. Sanitary and demographic indicators of Surkhandarya region, data on children's health, medical services for population of separate districts, statistics on condition of population of Surkhandarya region The



causes of helminthiasis in children were studied.

The cities of Termez, Denau, Shurchi, Altynsay, Boysun, Jarkurgan and Termez in Surkhandarya province were exposed to helminthiasis using the "incident monitoring" research method. Healthy children (300 children) were included in the control group, children with helminthiasis (300 children) were included in the incident group and the study was based on these groups and we studied the study in 4 stages. According to statistics, as of December 31, 2019 in Surkhandarya region there are 13 districts and 1 city in which 83.4 % of children registered with enterobiasis, ascariasis, hymenolepidosis at children aged till 14 years live in Termez, Denau, Shurchi, Altynsay, Boysun, Jarkurgan and Termez districts.

According to Table 1, if we look at the dynamics of helminths in children in the Surkhandarya region in 2018-2019, we can see a trend of increasing helminths in children. The helminth disease incidence in children increased slightly compared to 2018, including 6.8% by 300 children in 2018 and 7.3% by 2019.

The following helminths are the main causes of helminthiasis in children:

Askarid - when playing in the mud, leaving the street, do not wash your hands thoroughly with soap before eating and after going to the toilet, do not make sure to litter a person (often do not wash the toilet seats), use a few children without disinfected toys, do not wash fruits and vegetables completely (without cleaning or in raw form) when eating nails are not removed often, personal hygiene rules are not observed;

Striped worms - from well cooked or raw meat (worm eggs to pork, beef and fish);

Oysters may spread through toilet seats or when hygiene is not practiced, and when a person with oysters breathes, ostrich eggs are thrown into the air while breathing.

According to our study, enterobiasis in children under 14 years old in Surkhandarya region in 2018-2019 is 11.7% in Denau district, 10.9% in Jarkurgan district, 9.5% in Shurchi district, 9.1% in Boysun district, 8.9% in Sherabad district, 8.6% in Altynsay district, 8.1% in Termez district and 33.2% in other districts.

The prevalence of ascariasis by sex was 50.7% among boys in 2018, 49.3% among girls, 51% among boys in 2019, 49% among girls, 50.8% among boys in 2020 and 49.2% among girls.

This table shows that the prevalence of ascariasis by sex is slightly higher in boys than in girls.

Conclusion. According to our study, the incidence of enterobiasis in children in Surkhandarya region is 44% and that of ascariasis is 23%, which is the main cause of helminthiasis. Among all infected helminthoses, 78.9% of cases of enterobiasis in children, 71% of cases of ascariasis, 69.7% of ostrich, 58.5% of hymenolepidosis and 55% of hyardiasis in children.

Studies have shown that the most important requirement in the fight against helminthiasis is the prevention and choice of the right and effective medicines. In order to understand which drugs to use, it is necessary to undergo a thorough examination, determine the stage of the disease and only then begin treatment. During the prophylaxis, it is necessary to



monitor the sanitary condition of living, working and training areas. Fish and meat should be well roasted, as they may contain helminths. Immunity measures should be

taken. Effective treatment is not possible without proper understanding of adaptation mechanisms.

## References:

1. Sanitary epidemiological Surkhandarya region peace center report for 2017-2020.
2. Myandina I.G., Tarasenko V.E. Medical parasitology. Practical Medicine – 2013y. - 280 p.
3. Medical parasitology; Medicine - Moscow, 2012 y. - 304 p.
4. Jalilov J. J. Improving the prevention, care and social assistance of children with disabilities // Monograph. Tashkent-2020.
4. Kornakova EE. Medical parasitology. M.: Academy, 2010 y.-34p.
5. Jalilov J. J. The role of medical workers and scringing centers in preventing the birth of children with disabilities in the Surkhandarya region; International scientific journal; New day in medicine - 2019 y.-28-32 p.
6. Jalilov J. J. Indicators and causes of helminthiasis in children of surkhandarya region // Xorazim Ma'mun akademiyasi axborotnomasi, dekabr 2020y. 6-8 6.
7. Jalilov J. J. Causes and indicators of child disability in Surkhandarya region in 2017-2020 // Polish Science Journal", Poland, 2020. P. 116-119.
8. J.J. Jalilov, M.A. Saitmuratov, S.E. Xudoyberdiev, B.F. Danayev Characteristics of Child Helminthiasis in Surkhandarya Region // International Journal of Development and Public Policy. T-1. №4. P. 139-142.