



FUNCTION OF SPHINCTERS IN THE BODY

Djuraeva Barno Gulomovna

Abdujabborova Khurshida

Bakhodirova Mubina

<https://www.doi.org/10.5281/zenodo.10431810>

ARTICLE INFO

Received: 18th December 2023

Accepted: 25th December 2023

Online: 26th December 2023

KEY WORDS

Sphincters, Circular Muscles, Gatekeepers, Anatomy of Sphincters, Types of Sphincters, Muscular Sphincters, Smooth Muscle Sphincters, Striated Muscle Sphincters, Gastrointestinal Sphincters, Urinary Sphincters, Reproductive Sphincters, Cardiovascular Sphincters

ABSTRACT

This informative article delves into the multifaceted functions of sphincters in the human body. Acting as dynamic gatekeepers, sphincters are circular muscles strategically placed in various organs and systems to regulate the controlled flow of substances. The article explores the anatomy and different types of sphincters, ranging from muscular and smooth muscle sphincters to striated muscle sphincters, each serving unique roles. It delves into specific sphincters within the gastrointestinal, urinary, reproductive, cardiovascular, and respiratory systems, elucidating their roles in maintaining homeostasis. The article also addresses the crucial role of sphincters in preventing disorders and dysfunctions, emphasizing their significance in clinical interventions. By providing a comprehensive overview, this article serves as a valuable resource for understanding the intricate functions of sphincters in the human body.

Sphincters, often referred to as the body's gatekeepers, play a pivotal role in regulating the flow of substances within various organs and systems. These ring-like muscles act as dynamic barriers, allowing precise control over the movement of fluids and materials. This article delves into the diverse functions of sphincters throughout the human body, exploring their anatomy, types, and crucial roles in maintaining homeostasis.

Anatomy of Sphincters:

Sphincters are circular muscles that encircle an opening or passage in the body. They are strategically positioned at points where controlled passage is essential, serving as both protective guards and active regulators.

Types of Sphincters:

Muscular Sphincters: Found in the digestive system, these sphincters are primarily composed of smooth muscle and regulate the movement of materials through the digestive tract. Notable examples include the lower esophageal sphincter (LES) and pyloric sphincter.



Smooth Muscle Sphincters: Located in various organs, such as blood vessels and the respiratory system, smooth muscle sphincters control blood flow and airflow, ensuring optimal organ function.

Striated Muscle Sphincters: Voluntarily controlled, these sphincters are present in structures like the anus and urethra, allowing conscious control over the release of waste and urine.

Gastrointestinal Sphincters:

The digestive system hosts several sphincters crucial for maintaining the one-way flow of ingested materials. The LES prevents stomach acid from entering the esophagus, while the pyloric sphincter regulates the release of partially digested food into the small intestine.

Urinary and Reproductive Sphincters:

In the urinary system, the internal and external urethral sphincters control the release of urine from the bladder. Similarly, reproductive organs, such as the uterus, feature sphincters that aid in various functions, including menstruation and childbirth.

Cardiovascular Sphincters:

Within the cardiovascular system, sphincters in blood vessels regulate blood flow. The precapillary sphincters, for instance, control blood supply to capillaries, ensuring efficient nutrient and oxygen exchange in tissues.

Respiratory Sphincters:

In the respiratory system, sphincters in the bronchi help control airflow. These structures ensure proper ventilation and prevent the backflow of air, contributing to efficient gas exchange in the lungs.

Role in Homeostasis:

Sphincters are integral to maintaining homeostasis by precisely controlling the passage of substances. Whether regulating the movement of food, air, urine, or blood, sphincters contribute to the equilibrium required for optimal organ function.

Disorders and Dysfunctions: Malfunctions in sphincters can lead to various health issues. Conditions like gastroesophageal reflux disease (GERD), urinary incontinence, and certain cardiovascular disorders may involve impaired sphincter function.

Clinical Interventions:

Medical interventions often target sphincters to manage disorders. Surgical procedures, medications, and lifestyle modifications aim to restore or enhance sphincter function, addressing issues related to their malfunction.

To define sphincters, let us understand that these are special circular muscles able to open and close certain body parts. One of the vital sphincters [muscle](#) is the sphincter pylori, a thick middle layer of the stomach muscle opening to pylorus caving into the small intestine. Mostly, the sphincters muscle acts as a valve to regulate fluid such as bile, [urine](#), or faecal substance. The mechanism of the sphincter may be involuntary depending on the autonomic nervous system or voluntary operated by the somatic nervous system. The sphincter ani externus control the anal opening by contracting.

The sphincter urethrae are the most paramount voluntary mechanism of urination. Sphincter papillae are located in the iris, making it contract in the presence of bright luminosity.



The sphincter meaning is a ring-shaped muscle that relaxes or contracts to open or close a body passage or opening. To define sphincter, these muscles are smooth muscles that remain in a contraction state. These are specialized latch systems in the contractile filaments permitting sphincters to maintain contracting position with minimum effort.

The internal anal sphincter is an exception that blocks the backward flow of intraluminal contents. The lower oesophageal sphincter thwarts gastric acid into the oesophagus. Dysfunction of the oesophageal sphincter causes continuous exposure of oesophageal mucosa to gastric acid, which can cause dysplastic heartburn that may become cancerous. The sphincters muscle function of external anal sphincters is to control defecation; it is a short tube of skeletal muscle surrounding the inferior portion of the anal canal.

Sphincters Muscle Stomach

There are six different sphincter muscles in the abdomen. The upper oesophageal sphincter (UES) is located at the end of the pharynx, where it guards the opening of the oesophagus. UES prevents the induction of air in the oesophagus while breathing and prevents food from getting inside the respiratory system. Because of the location in the pharynx, it also controls vomiting and burping.

The lower oesophageal sphincter (LES), also called cardiac sphincter, is located at the end of the oesophagus, where it joins the stomach. The principal function of LES is to permit food to pass from the oesophagus to the stomach and to enable air to pass from the stomach while burping. It also prevents the backward flow of gastric acid into the oesophagus.

Pyloric Sphincter. The pyloric sphincter is situated between the stomach and duodenum. This sphincter allows chyme (partially digested food) to transmit from the stomach to the duodenum for further digestion and absorption of nutrients through mucus into the blood.

Sphincter of Oddi. The sphincter of Oddi (SO) is located at the influx of the bile duct and pancreatic duct at the duodenum. SO opens to allow bile from the gallbladder and enzymes from the pancreas to mix to break down food particles for assimilation.

Ileocecal Sphincter. At the junction of the small and large intestine, the ileocecal sphincter is situated. The functionality of this sphincter is not clear, except it allows chyme from the small intestine to the large intestine.

Anal Sphincter. This sphincter is located at the end of the gastrointestinal tract at the last part of the rectum. This muscle has both inner and outer components to allow the passage of the stool. The automatic nervous system controls the inner sphincter while the outer is managed by the somatic nervous system. There are other sphincters in the body as well.

What is the Function of Sphincter Muscles?

The urethral sphincter controls the holding and pouring of urine. Similar to the anal sphincter, it has inner and outer muscles controlled by the involuntary and voluntary nervous system.

Iris Sphincter. This sphincter, also known as sphincter papilla, is located at the iris, the coloured part of the eye. It surrounds the pupil of the iris and contracts the pupil in bright light through the pupillary light reflex or during accommodation. The diameter of the pupil controls the amount of light that enters the iris reaching at the back of the eye at the retina.

Sphincter muscles examples are iris sphincter, anal sphincter, ileocecal sphincter, pyloric sphincter; there are about fifty to sixty different types of sphincter muscles in the



body. Some are microscopic, such as millions of precapillary sphincters in the circulatory system. Some are controlled by the involuntary nervous system; some respond to specific stimuli; others are controlled by the voluntary nervous system. Sphincter muscles can degenerate or be damaged, leading to several health issues. The dysfunction of the lower oesophageal sphincter (LES) permits gastro acid to enter the oesophagus resulting in acid reflux.

Conclusion: In essence, sphincters emerge as crucial players in the symphony of bodily functions. Their intricate coordination ensures the controlled movement of substances, safeguarding organs from potential harm. This article offers a glimpse into the diverse functions of sphincters, underscoring their significance in maintaining the delicate balance required for the human body's optimal performance.

References:

1. Atkinson, M., Edwards, D. A. W., Honour, A. J. and Rowlands, E. N. (1957). The oesophagogastric sphincter in hiatus hernia. *Lancet*, ii, 1138
2. Bargaen, J. A., Wesson, H. R. and Jackman, R. J. (1940). Studies on the ileocaecal junction (ileocecus). *Surg., Gynec. Obst.*, 71, 33
3. Beattie, J. (1924). The early stages of the development of the ileocolic sphincter. *J. Anat.*, 59, 56
4. Boyden, E. A. (1937). The sphincter of Oddi in man and certain representative mammals. *Surgery*, 1, 25
5. Colp. R., Doubilet, H. and Gerber, I. E. (1936). Endocholedochal section of the sphincter of Oddi. *Arch. Surg.*, 33, 696
6. Courtney, H. (1950). Anatomy of the pelvic diaphragm and anorectal musculature as related to sphincter preservation in anorectal surgery. *Am. J. Surg.*, 79, 155
7. Fleischner, F. G. and Bernstein, C. (1950). Roentgen-anatomical studies of the normal ileocaecal valve. *Radiology*, 54, 43
8. Goligher, J. C., Leacock, A. G. and Brossy, J. J. (1955). The surgical anatomy of the anal canal. *Br. J. Surg.*, 43, 51
9. Horton, B. T. (1928). Pyloric musculature, with special reference to pyloric block. *Am. J. Anat.*, 41, 197
10. Killian, G. (1908). Ueber den Mund der Speiseröhre. *Zeitschr. Ohrenh.*, 55, 1
11. Kirk, J. (1944). Observations on the histology of the choledochoduodenal junction and papilla duodeni, with particular reference to the ampulla of Vater and sphincter of Oddi. *J. Anat.*, 78, 118
12. Kreilkamp, B. L. and Boyden, E. A. (1940). Variability in the composition of the sphincter of Oddi: a possible factor in the pathologic physiology of the biliary tract. *Anat. Rec.*, 76, 485
13. Lendrum, F. C. (1937). Anatomic features of the cardiac orifice of the stomach: with special reference to cardiospasm. *Arch. Int. Med.*, 59, 474
14. Levy, E. (1936). Anorectal musculature. *Am. J. Surg.*, 34, 141
15. Milligan, E. T. C. and Morgan, C. N. (1934). Surgical anatomy of the anal canal, with special reference to anorectal fistulae. *Lancet*, ii, 1150, 1213



16. Moersch, H. J. and Judd, E. S. (1934). Diagnosis and treatment of pharyngo-oesophageal diverticulum. *Surg., Gynec. Obst.*, 58, 781
17. Morgan, C. N. (1949). The surgical anatomy of the ischio-rectal space. *Proc. R. Soc. Med.*, 42, 189
18. Moynihan, B. (1927). Diverticula of the alimentary canal. *Lancet*, i, 1061
19. Puestow, C. B. (1938). Changes in intra-choledochal pressure following cholecystectomy. *Surg., Gynec. Obst.*, 67, 82
20. Ramsay, E. H., Watson, J. S., Gramiak, R. and Weinberg, S. A. (1955). Cinefluorographic analysis of the mechanism of swallowing. *Radiology*, 64, 498
21. Rosenberg, J. C. and Didio, L. J. (1969). In vivo appearance and function of the ileum as observed directly through a cecostomy. *Am. J. Gastroenterol.*, 52, 411