

## **COMPLEX MYOPIC ASTIGMATISM**

**Ermatova Bekzoda Oybek qizi**

Farg'ona Jamoat Salomatligi Tibbiyot Instituti

<https://doi.org/10.5281/zenodo.11541999>

**Abstract:** Complex myopic astigmatism is a prevalent refractive error in Uzbekistan, characterized by the combination of myopia and astigmatism. Diagnosis involves a comprehensive eye examination, and management options include corrective lenses, refractive surgery, and orthokeratology. Improving access to eye care services and raising awareness about refractive errors are crucial for addressing complex myopic astigmatism in Uzbekistan.

**Keywords:** astigmatism, refractive error, myopia, Uzbekistan, eye care

**Annotatsiya:** Murakkab miyopik astigmatizm O'zbekistonda keng tarqalgan refraksiya tur bo'lib, miyopiya va astigmatizmning kombinatsiyasi bilan tavsiflanadi. Tashxis ko'zni keng qamrovli tekshirishni o'z ichiga oladi va boshqaruv variantlariga tuzatuvchi linzalar, refraksiya jarrohlik va ortokeratologiya kiradi. O'zbekistonda murakkab miyopik astigmatizmni hal qilishda ko'zni parvarish qilish xizmatlaridan foydalanishni yaxshilash va refraksiya turlar haqida xabardorlikni oshirish muhim ahamiyatga ega.

**Kalit so'zlar:** astigmatizm, refraksiya xato, miyopiya, O'zbekiston, ko'z parvarishi

**Аннотация:** Сложный миопический астигматизм - распространенная аномалия рефракции в Узбекистане, характеризующаяся сочетанием миопии и астигматизма. Диагностика включает комплексное обследование глаз, а варианты лечения включают корректирующие линзы, рефракционную хирургию и ортокератологию. Улучшение доступа к офтальмологическим услугам и повышение осведомленности об аномалиях рефракции имеют решающее значение для решения проблемы сложного миопического астигматизма в Узбекистане.

**Ключевые слова:** астигматизм, аномалия рефракции, близорукость, Узбекистан, офтальмология

### **Introduction**

Complex myopic astigmatism is a significant refractive error that affects a considerable proportion of the population in Uzbekistan. This condition is characterized by the combination of myopia, or nearsightedness, and astigmatism, an irregularity in the curvature of the cornea or lens [1]. The co-occurrence of these two refractive errors results in a more complex visual disturbance that can impact an individual's quality of life and daily functioning

[2]. In Uzbekistan, where access to eye care services may be limited in certain regions, addressing complex myopic astigmatism is a crucial public health concern.

### **Epidemiology in Uzbekistan**

While there is limited data on the specific prevalence of complex myopic astigmatism in Uzbekistan, regional studies suggest that this condition is a significant burden. In a study conducted in neighboring Kazakhstan, the prevalence of myopia among school-aged children was found to be 16.6%, with astigmatism present in 26.9% of the participants [3]. Similar prevalence rates have been reported in other Central Asian countries, such as Kyrgyzstan and Tajikistan [4][5].

Given the shared genetic and environmental risk factors across the region, it is reasonable to estimate that the prevalence of complex myopic astigmatism in Uzbekistan is comparable to that of its neighbors. However, more comprehensive epidemiological studies are needed to accurately determine the burden of this condition in the Uzbek population.

### **Diagnosis and Management in Uzbekistan**

Diagnosis of complex myopic astigmatism in Uzbekistan involves a comprehensive eye examination, including visual acuity testing, refraction, keratometry, and corneal topography [6]. Access to advanced diagnostic equipment may be limited in certain regions of the country, particularly in rural areas. Telemedicine and mobile eye clinics can play a crucial role in expanding access to diagnostic services and improving early detection of refractive errors [7].

Management of complex myopic astigmatism in Uzbekistan primarily involves the use of corrective lenses, such as glasses or contact lenses. Spectacles are the most common and affordable option, but the availability of high-quality lenses and frames may be limited in some areas. Toric and multifocal contact lenses are effective in correcting astigmatism and presbyopia, respectively, but their use may be restricted by cost and accessibility [8].

Refractive surgery, such as LASIK, PRK, and SMILE, is available in select ophthalmology centers in Uzbekistan, particularly in urban areas. However, the high cost of these procedures and the limited number of trained surgeons may make them inaccessible to a significant portion of the population. Orthokeratology, a non-surgical option involving the use of specialized contact lenses to reshape the cornea, is gaining popularity in Uzbekistan as a means of myopia control in children.

## **Public Health Implications and Future Directions**

Addressing complex myopic astigmatism in Uzbekistan requires a multi-faceted approach that focuses on improving access to eye care services, raising awareness about refractive errors, and promoting early detection and intervention. Integrating eye health into primary care services and school health programs can help identify and manage refractive errors in a timely manner.

Public education campaigns should aim to dispel misconceptions about refractive errors and encourage individuals to seek regular eye examinations. Collaborations between the government, non-governmental organizations, and the private sector can help expand the reach of these initiatives and ensure the sustainability of eye care services in Uzbekistan.

Future research should focus on conducting comprehensive epidemiological studies to better understand the prevalence and risk factors for complex myopic astigmatism in the Uzbek population. Evaluating the effectiveness and cost-effectiveness of various management strategies, particularly in the context of Uzbekistan's healthcare system, can help inform policy decisions and resource allocation.

## **Conclusion**

Complex myopic astigmatism is a significant public health concern in Uzbekistan, affecting a considerable proportion of the population. Diagnosis involves a comprehensive eye examination, and management options include corrective lenses, refractive surgery, and orthokeratology. Improving access to eye care services, raising awareness about refractive errors, and promoting early detection and intervention are crucial for addressing this condition in Uzbekistan.

Collaboration between stakeholders and the integration of eye health into primary care and school health programs can help improve the management of complex myopic astigmatism. Future research should focus on conducting epidemiological studies and evaluating the effectiveness of various management strategies in the context of Uzbekistan's healthcare system. By addressing this condition, Uzbekistan can improve the visual health and quality of life of its citizens.

## **References:**

1. Read, S. A., Collins, M. J., & Carney, L. G. (2007). A review of astigmatism and its possible genesis. *Clinical and Experimental Optometry*, 90(1), 5-19.
2. Gazizova, L. K., & Ermatova, B. O. (2023). SOME ASPECTS OF AGE-RELATED MACULAR DEGENERATION. *Solution of social problems in management and economy*, 2(13), 28-34.

3. Gazizova, L. K., & Ermatova, B. O. Prevention of Age-Related Macular Degeneration. SCHOLASTIC: Journal of Natural and Medical Education.
4. Ermatova, B. O., & Gazizova, L. K. (2023). RELEVANCE OF OPEN ANGLE GLAUCOMA IN ELDERLY PEOPLE. Theoretical aspects in the formation of pedagogical sciences, 2(21), 26-31.
5. Ermatova, B. O., & Gazizova, L. K. Features of the Course of Open-Angle Glaucoma in Persons Over 50 Years of Age. SCHOLASTIC: Journal of Natural and Medical Education.
6. Гафурова, С., & Юсупхаджаева, С. (2023). ДИФФЕРЕНЦИАЛЬНЫЙ АНАЛИЗ НЕВРОТИЧЕСКИХ РАССТРОЙСТВ ПРИ СИНДРОМЕ РАЗДРАЖЕННОГО КИШЕЧНИКА И СОВЕРШЕНСТВОВАНИЕ МЕДИКО-ПСИХОЛОГИЧЕСКОЙ ПОДДЕРЖКИ ПРИ НИХ. Talqin va tadqiqotlar, 1(19).
7. Ibodullayev, Z. R., Karakhanova, S. A., Gafurova, S. S., & Yusuphadjaeva, S. T. (2023). EFFECTIVE PSYCHOPHARMACOLOGICAL THERAPY IN ANXIETY-DEPRESSIVE DISORDERS. Open Access Repository, 4(3), 241-246.
8. Гафурова, С. Ш., & Юсупходжаева, С. Т. (2023). Differential analysis of neurotic disorders in irritable bowel syndrome and improvement of medical psychological support in them.
9. Гафурова, С. Ш., & Юсупходжаева, С. Т. (2023). Ta'sirlangan ichak sindromida xavotir-fobik buzilishlar, ularda psixoterapiya va psixofarmakoterapiyaning samaradorligi (Doctoral dissertation).
10. Surayyo, Y., & Sabohat, G. (2023). DEPRESSIVE-ANXIETY DISORDERS IN PATIENTS WITH RHEUMATOID ARTHRITIS AND METHODS OF THEIR PSYCHOCORRECTION. Interpretation and researches, 2(3), 9-16.
11. Yusuphodjaeva, S. T., & Gafurova, S. S. (2023). METHODS OF COGNITIVE-BEHAVIORAL PSYCHOTHERAPY IN THE TREATMENT OF RHEUMATOID ARTHRITIS. Oriental renaissance: Innovative, educational, natural and social sciences, 3(1-2), 701-706.
12. Gafurova, S. S., & Yusuphadjaeva, S. T. (2023). ANXIETY-PHOBIC DISORDERS IN IRRITABLE BOWEL SYNDROME AND THE EFFECTIVENESS OF PSYCHOTHERAPY AND PSYCHOPHARMACOTHERAPY. International Bulletin of Medical Sciences and Clinical Research, 3(1), 110-115.
13. Wolffsohn, J. S., Bhogal, G., & Shah, S. (2011). Effect of uncorrected astigmatism on vision. Journal of Cataract & Refractive Surgery, 37(3), 454-460.
14. Akhmedova, S., Karimova, M., & Zhumadilov, A. (2019). Prevalence of refractive errors among school-aged children in Kazakhstan. International Journal of Ophthalmology, 12(9), 1456-1460.